

ON TWO FORMULAS CONTIGUOUS TO A TRANSFORMATION DUE TO BAILEY

NEETHU. P, MEDHAT A. RAKHA¹ AND ARJUN K. RATHIE

ABSTRACT. The aim of this short research note is to provide two interesting formulas contiguous to the following transformation due to Bailey (written here in corrected form) viz.

$$\begin{aligned} & (1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; \\ 2b + \frac{1}{2}; & \end{matrix} -\frac{4x}{(1-x)^2} \right] \\ &= {}_2F_1 \left[\begin{matrix} 1, & \frac{3}{4} - b; \\ \frac{3}{4} + b; & x^2 \end{matrix} \right] - \frac{x(4b-1)}{(4b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{5}{4} - b; \\ \frac{5}{4} + b; & x^2 \end{matrix} \right] \end{aligned}$$

The results are derived with the help of two formulas contiguous to classical Whipple's summation theorem on the sum of a ${}_3F_2$ obtained earlier by Lavoie, et al.

The results established in this short research note are simple, interesting, easily established and may be useful.

1. INTRODUCTION

In the theory of hypergeometric and generalized hypergeometric series, classical summation theorems such as those of Gauss, Gauss second, Kummer and Bailey for the series ${}_2F_1$; Watson, Dixon, Whipple and Saalschütz for the series ${}_3F_2$ and others play an important role. Several formulae were given by Gauss and Kummer expressing the product of two hypergeometric series, such as $e^{-x} {}_1F_1(x)$ as a series of the type ${}_1F_1(-x)$ and

$$(1+x)^{-p} {}_2F_1 \left(\frac{4x}{(1+x)^2} \right)$$

as a series of the type ${}_2F_1(x)$. By employing the above mentioned classical summation theorems, in 1928, Bailey [1] made a systematic search for several formulae. Evidently if the product of two hypergeometric series can be expressed as a hypergeometric series with argument x , the coefficient of x^n in the product must be expressible in terms of the Gamma function.

¹corresponding author

2010 *Mathematics Subject Classification.* 33C05, 33C20, 33C70.

Key words and phrases. Generalized hypergeometric functions, Bailey's transformation, Whipple's summation theorem.

Recently good progress has been done in the direction of generalizing and extending the above mentioned classical summation theorems. For this, we refer to the research papers [2] - [5].

In our present investigation, we are interested in the following transformation due to Bailey [1]

$$(1.1) \quad (1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; \\ 2b + \frac{1}{2}; \end{matrix} \quad -\frac{4x}{(1-x)^2} \right] = {}_2F_1 \left[\begin{matrix} 1, & \frac{3}{4} - b; \\ \frac{3}{4} + b; \end{matrix} \quad x^2 \right] - \frac{x(4b-1)}{(4b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{5}{4} - b; \\ \frac{5}{4} + b; \end{matrix} \quad x^2 \right].$$

Bailey [1] obtained this result by employing the following classical Whipple's summation theorem [8] viz

$$(1.2) \quad {}_3F_2 \left[\begin{matrix} a, & b, & c; \\ & e, & f; \end{matrix} \quad 1 \right] = \frac{\pi \Gamma(e) \Gamma(f)}{2^{2c-1} \Gamma(\frac{1}{2}a + \frac{1}{2}e) \Gamma(\frac{1}{2}a + \frac{1}{2}f) \Gamma(\frac{1}{2}b + \frac{1}{2}e) \Gamma(\frac{1}{2}b + \frac{1}{2}f)}$$

provided $a + b = 1$ and $e + f = 1 + 2c$.

In 1996, Lavoie, et al. [5] generalized the above mentioned classical Whipple's summation theorem (1.2) and have obtained explicit expressions of

$$(1.3) \quad {}_3F_2 \left[\begin{matrix} a, & b, & c; \\ & e, & f; \end{matrix} \quad 1 \right]$$

provided $a + b = 1 + i + j$ and $e + f = 2c + i + 1$ for $i, j = 0, \pm 1, \pm 2, \pm 3$.

In our present investigation, we shall require the following results which can be obtained from (1.3) by taking $i = 1, j = -1$ and $i = -1, j = 1$ respectively, these are

- For $i = 1, j = -1$

$$(1.4) \quad {}_3F_2 \left[\begin{matrix} a, & b, & c; \\ & e, & f; \end{matrix} \quad 1 \right] = \frac{\Gamma(e) \Gamma(f) \Gamma(e-c-1)}{2^{2a} \Gamma(e-a) \Gamma(f-a) \Gamma(e-c)} \times \left\{ \frac{\Gamma(\frac{e}{2} - \frac{a}{2} + \frac{1}{2}) \Gamma(\frac{f}{2} - \frac{a}{2})}{\Gamma(\frac{e}{2} + \frac{a}{2} - \frac{1}{2}) \Gamma(\frac{f}{2} + \frac{a}{2})} - \frac{\Gamma(\frac{e}{2} - \frac{a}{2}) \Gamma(\frac{f}{2} - \frac{a}{2} + \frac{1}{2})}{\Gamma(\frac{e}{2} + \frac{a}{2}) \Gamma(\frac{f}{2} + \frac{a}{2} - \frac{1}{2})} \right\}$$

provided $a + b = 1, e + f = 2c + 2$.

- For $i = -1, j = 1$

$$(1.5) \quad {}_3F_2 \left[\begin{matrix} a, & b, & c; & \\ & e, & f; & 1 \end{matrix} \right] = \frac{\Gamma(e) \Gamma(f) \Gamma(c-1)}{2^{2a} \Gamma(e-a) \Gamma(f-a) \Gamma(c)} \times \left\{ \frac{\Gamma\left(\frac{e}{2} - \frac{a}{2} + \frac{1}{2}\right) \Gamma\left(\frac{f}{2} - \frac{a}{2}\right)}{\Gamma\left(\frac{e}{2} + \frac{a}{2} - \frac{1}{2}\right) \Gamma\left(\frac{f}{2} + \frac{a}{2}\right)} + \frac{\Gamma\left(\frac{e}{2} - \frac{a}{2}\right) \Gamma\left(\frac{f}{2} - \frac{a}{2} + \frac{1}{2}\right)}{\Gamma\left(\frac{e}{2} + \frac{a}{2}\right) \Gamma\left(\frac{f}{2} + \frac{a}{2} - \frac{1}{2}\right)} \right\}$$

provided $a + b = 1, e + f = 2c$.

The aim of this short research note is to find the explicit expressions of

$$(1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; & \\ & 2b + \frac{1}{2} + l; & -\frac{4x}{(1-x)^2} \end{matrix} \right]$$

for $l = \pm 1$.

The results are derived with the help of contiguous Whipple's summation theorems (1.4) and (1.5). The results established in this short research note are simple interesting, easily established and may be useful.

2. MAIN RESULTS

The transformation formulas closely related to the Bailey's transformation to be established in this short research note are given by

$$(2.1) \quad \begin{aligned} & (1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; & \\ & 2b + \frac{3}{2}; & -\frac{4x}{(1-x)^2} \end{matrix} \right] \\ &= \frac{1}{2(2b+1)} {}_3F_2 \left[\begin{matrix} 1, & \frac{5}{4}, & \frac{1}{4} - b; & \\ \frac{1}{4}, & \frac{5}{4} + b; & & x^2 \end{matrix} \right] + \frac{(4b+1)}{2(2b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{3}{4} - b; & \\ \frac{3}{4} + b; & & x^2 \end{matrix} \right] \\ &+ \frac{3(4b+1)x}{2(2b+1)(4b+3)} {}_3F_2 \left[\begin{matrix} 1, & \frac{7}{4}, & \frac{3}{4} - b; & \\ \frac{3}{4}, & \frac{7}{4} + b; & & x^2 \end{matrix} \right] - \frac{(4b-1)x}{2(2b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{5}{4} - b; & \\ \frac{5}{4} + b; & & x^2 \end{matrix} \right] \end{aligned}$$

and

$$\begin{aligned}
 & (1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; \\ 2b - \frac{1}{2}; & -\frac{4x}{(1-x)^2} \end{matrix} \right] \\
 &= \frac{1}{4(1-b)} {}_3F_2 \left[\begin{matrix} 1, & \frac{5}{4}, & \frac{5}{4} - b; \\ \frac{1}{4}, & b + \frac{1}{4}; & x^2 \end{matrix} \right] - \frac{(4b+3)}{4(1-b)} {}_2F_1 \left[\begin{matrix} 1, & \frac{7}{4} - b; \\ b - \frac{1}{4}; & x^2 \end{matrix} \right] \\
 (2.2) \quad & + \frac{3(4b-3)x}{4(1-b)(4b-1)} {}_3F_2 \left[\begin{matrix} 1, & \frac{7}{4}, & \frac{7}{4} - b; \\ \frac{3}{4}, & \frac{3}{4} + b; & x^2 \end{matrix} \right] + \frac{(4b-5)x}{4(1-b)} {}_2F_1 \left[\begin{matrix} 1, & \frac{9}{4} - b; \\ b + \frac{1}{4}; & x^2 \end{matrix} \right]
 \end{aligned}$$

Proof. In order to derive (2.1), we proceed as follows. Denoting the left-hand side of (2.1) by S , we have

$$S = (1-x)^{-1} {}_2F_1 \left[\begin{matrix} 1, & b; \\ 2b + \frac{3}{2}; & -\frac{4x}{(1-x)^2} \end{matrix} \right].$$

Expressing ${}_2F_1$ as a series, we have after some algebra

$$S = \sum_{n=0}^{\infty} \frac{(b)_n (-1)^n 2^{2n}}{(2b + \frac{3}{2})_n} x^n (1-x)^{-(2n+1)},$$

using Binomial theorem, we have

$$S = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(b)_n (-1)^n 2^{2n}}{(2b + \frac{3}{2})_n} \frac{(2n+1)_m}{m!} x^{n+m}.$$

Now, replacing m by $m - n$ and using the known result [6, Lemma 10, p. 56 Eq. (1)]

$$\sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A(k, n) = \sum_{n=0}^{\infty} \sum_{k=0}^n A(k, n-k),$$

we have

$$S = \sum_{m=0}^{\infty} \sum_{n=0}^m \frac{(b)_n (-1)^n 2^{2n}}{(2b + \frac{3}{2})_n} \frac{(2n+1)_{m-n}}{(m-n)!} x^m,$$

using the elementary identities [6]

$$\begin{aligned}
 (a)_{m-n} &= \frac{(-1)^n (a)_m}{(1-a-m)_n}, \\
 (m-n)! &= \frac{(-1)^n m!}{(-m)_n}
 \end{aligned}$$

and

$$\Gamma(2z) = \frac{2^{2z-1} \Gamma(z) \Gamma(z + \frac{1}{2})}{\sqrt{\pi}},$$

we have, after some simplification

$$S = \sum_{m=0}^{\infty} x^m \sum_{n=0}^m \frac{(-m)_n (1+m)_n (b)_n}{(\frac{1}{2})_n (2b + \frac{3}{2})_n n!}.$$

Summing up the inner series, we have

$$S = \sum_{m=0}^{\infty} x^m {}_3F_2 \left[\begin{matrix} -m, & 1+m, & b; \\ \frac{1}{2}, & 2b + \frac{3}{2}; \end{matrix} \quad 1 \right].$$

Now, separating into even and odd powers of x , we have

$$\begin{aligned} S &= \sum_{m=0}^{\infty} x^{2m} {}_3F_2 \left[\begin{matrix} -2m, & 1+2m, & b; \\ \frac{1}{2}, & 2b + \frac{3}{2}; \end{matrix} \quad 1 \right] \\ &+ x \sum_{m=0}^{\infty} x^{2m} {}_3F_2 \left[\begin{matrix} -2m-1, & 2m+2, & b; \\ \frac{1}{2}, & 2b + \frac{3}{2}; \end{matrix} \quad 1 \right]. \end{aligned}$$

Now assume that

$$(2.3) \quad S = A + B,$$

then we have

$$A = \sum_{m=0}^{\infty} x^{2m} {}_3F_2 \left[\begin{matrix} -2m, & 1+2m, & b; \\ \frac{1}{2}, & 2b + \frac{3}{2}; \end{matrix} \quad 1 \right].$$

It is now, easy to see that, the ${}_3F_2$ can be evaluated with the help of the contiguous Whipple's summation theorem (1.4) by taking $a = -2m, b = 1 + 2m, c = b, e = \frac{1}{2}$ and $f = 2b + \frac{3}{2}$ and after some simplification, we get

$$\begin{aligned} A &= \frac{1}{2(2b+1)} \sum_{m=0}^{\infty} \frac{(1)_m \left(\frac{5}{4}\right)_m \left(\frac{1}{4} - b\right)_m}{\left(\frac{1}{4}\right)_m \left(\frac{5}{4} + b\right)_m m!} x^{2m} \\ &+ \frac{(4b+1)}{2(2b+1)} \sum_{m=0}^{\infty} \frac{(1)_m \left(\frac{3}{4} - b\right)_m}{\left(\frac{3}{4} + b\right)_m m!} x^{2m} \end{aligned}$$

summing up the series, we have

$$\begin{aligned} (2.4) \quad A &= \frac{1}{2(2b+1)} {}_3F_2 \left[\begin{matrix} 1, & \frac{5}{4}, & \frac{1}{4} - b; \\ \frac{1}{4}, & b + \frac{5}{4}; \end{matrix} \quad x^2 \right] \\ &+ \frac{(4b+1)}{2(2b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{3}{4} - b; \\ \frac{3}{4} + b; \end{matrix} \quad x^2 \right]. \end{aligned}$$

In exactly the same manner, we can show that

$$\begin{aligned} (2.5) \quad B &= \frac{3(4b+1)x}{2(2b+1)(4b+3)} {}_3F_2 \left[\begin{matrix} 1, & \frac{7}{4}, & \frac{3}{4} - b; \\ \frac{3}{4}, & \frac{7}{4} + b; \end{matrix} \quad x^2 \right] \\ &- \frac{(4b-1)x}{2(2b+1)} {}_2F_1 \left[\begin{matrix} 1, & \frac{5}{4} - b; \\ \frac{5}{4} + b; \end{matrix} \quad x^2 \right]. \end{aligned}$$

Substituting the results of A and B from (2.4) and (2.5) in (2.3), we easily arrive at the right-hand side of our first main result (2.1). This completes the proof of (2.1).

In exactly the same manner, the result (2.2) can be established by employing the contiguous Whipple's summation theorem (1.5), so we prefer to omit the details. $\square$

Clearly the results (2.1) and (2.2) are closely related to the Bailey's transformation (1.1).

AUTHORS' CONTRIBUTIONS

The authors have equal contributions to each part of this paper. All authors have read and approved the final manuscript.

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

- [1] W. N. BAILEY: *Product of generalized hypergeometric series*, Proc. London Math. Soc. (2), 28 (1928), 242 – 254.
- [2] Y. S. KIM, M. A. RAKHA, A. K. RATHIE: *Extensions of classical summation theorems for the series ${}_2F_{1,3}F_2$ and ${}_4F_3$ with applications in Ramanujan's summations*, Int. J. Math. & Math. Sci., ID 309503 (2010), 26 pages.
- [3] J. L. LAVOIE, F. GRONDIN, A. K. RATHIE: *Generalizations of Watson's theorem on the sum of ${}_3F_2$* , Indian J. Math., 34 (1992), 23 – 32.
- [4] J. L. LAVOIE, F. GRONDIN, A. K. RATHIE, K. ARORA: *Generalizations of Dixon's theorem on the sum of ${}_3F_2$* , Math. Comp. American Math. Society, 62, No. 205 (1994), 267 – 276.
- [5] J. L. LAVOIE, F. GRONDIN, A. K. RATHIE: *Generalizations of Whipple's theorem on the sum of ${}_3F_2$* , J. Comput. Appl. Math., 72 (1996), 293 – 300.
- [6] E. D. RAINVILLE: *Special Functions*, The Macmillan Company, New York, 1960.
- [7] M. A. RAKHA, A. K. RATHIE: *Generalizations of classical summation theorems for the series ${}_2F_1$ and ${}_3F_2$ with applications*, Integral Transform and Special Functions, 22 (11) (2011) 823 – 840.
- [8] L. J. SLATER: *Generalized hypergeometric functions*, Cambridge University Press, Cambridge, 1966.

DEPARTMENT OF MATHEMATICS
SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES
CENTRAL UNIVERSITY OF KERALA, RIVERSIDE TRANSIT CAMPUS
PADENNAKKAD P.O. NILESHWAR, KASARAGOD - 671 328, KERALA - INDIA
E-mail address: neethukanayi82@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS
COLLEGE OF SCIENCE
SULTAN QABOOS UNIVERSITY
P.O.BOX 36 - AL-KHOUD 123
MUSCAT - SULTANATE OF OMAN
E-mail address: medhat@squ.edu.om

DEPARTMENT OF MATHEMATICS
SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES
CENTRAL UNIVERSITY OF KERALA, RIVERSIDE TRANSIT CAMPUS
PADENNAKKAD P.O. NILESHWAR, KASARAGOD - 671 328, KERALA - INDIA
E-mail address: akrathie@gmail.com